

NDUFS4 Antibody

CAB13519

Description

This NDUFS4 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB13519

Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: AQDQ, CI-18, MC1DN1, CI-AQDQ, CI-18 kDa, NDUFS4

Clone: -

Applications: **WB** **IHC-P** **IF/ICC** **ELISA**

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 4724

Uniprot: AB_2760380

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 20kDa

Calculated MW: 20kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: MCF7, HepG2, 293T, Mouse brain, Mouse bone marrow, Mouse liver, Rat brain, Rat liver


Recommended Dilutions:	WB	1:500 - 1:2000
	IHC-P	1:50 - 1:200
	IF/ICC	1:50 - 1:200
	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Western blot analysis of various lysates using NDUFS4 Rabbit pAb (CAB13519) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using NDUFS4 Rabbit pAb (CAB13519) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.