

RPL30 Antibody

CAB13690

Description

This RPL30 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB13690
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: L30, eL30, RPL30
Clone: -
Applications: **WB** | **IHC-P** | **IF/ICC** | **ELISA**
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat


Antibody Data

Gene ID: 6156
Uniprot: AB_2760550
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 13 kDa
Calculated MW: 13 kDa


Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.								
	Store Bradford Reagent at Room Temperature for 1 Year.								
Positive Sample:	HeLa, MCF7, HepG2, A-549, Mouse brain, Mouse heart, Mouse kidney, Rat liver								
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:1000 - 1:3000</td></tr> <tr> <td>IHC-P</td><td>1:50 - 1:100</td></tr> <tr> <td>IF/ICC</td><td>1:50 - 1:200</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:1000 - 1:3000	IHC-P	1:50 - 1:100	IF/ICC	1:50 - 1:200	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:1000 - 1:3000								
IHC-P	1:50 - 1:100								
IF/ICC	1:50 - 1:200								
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol								


Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB13690) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Rat heart using Rabbit pAb (CAB13690) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Confocal immunofluorescence analysis of HeLa cells using Rabbit pAb (CAB13690) at dilution of 1:200. Blue: DAPI for nuclear staining.