

Fukutin Monoclonal Antibody

CAB1385

Description

This Fukutin Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB1385

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Monoclonal Antibody

Synonyms: FCMD, CMD1X, LGMD2M, MDDGA4, MDDGB4, MDDGC4, LGMDR13, Fukutin

Clone: ARC2559

Applications: WB IHC-P ELISA

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 2218

Uniprot: -

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 54kDa

Calculated MW: 54kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: BxPC-3, RD, U-87MG, Mouse testis, Mouse brain, Mouse heart, Rat liver

Recommended Dilutions:


WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Western blot analysis of various lysates using Fukutin Rabbit mAb (CAB1385) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Human colon using Fukutin Rabbit mAb (CAB1385) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.