

TRIM9 Antibody

CAB13896

Description

This TRIM9 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB13896
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: RNF91, SPRING, TRIM9
Clone: -
Applications: **WB** **ELISA**
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat


Antibody Data

Gene ID: 114088
Uniprot: AB_2760750
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 79kDa
Calculated MW: 79kDa

Preparation & Storage

| | | | | | |
|---|--|-----------|----------------|--------------|---|
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. Store Bradford Reagent at Room Temperature for 1 Year. | | | | |
| Positive Sample: | HeLa, HepG2, DU145, Mouse brain, Rat brain | | | | |
| Recommended Dilutions: | <table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | WB | 1:500 - 1:2000 | ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| WB | 1:500 - 1:2000 | | | | |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. | | | | |
| Protein Quantification (Optional): | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol | | | | |

Validation Data



Western blot analysis of various lysates using TRIM9 Rabbit pAb (CAB13896) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.