

## SNAP23 Antibody

**CAB13909**

### Description

---

This SNAP23 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

|                      |   |
|----------------------|---|
| <b>SKU:</b>          | CAB13909  |
| <b>Contents:</b>     | 20 $\mu$ L, 100 $\mu$ L<br>Bradford Reagent: 1 vial (2ml) |
| <b>Category:</b>     | Polyclonal Antibody                                       |
| <b>Synonyms:</b>     | SNAP-23, SNAP23A, SNAP23B, HsT17016, SNAP23               |
| <b>Clone:</b>        | -   |
| <b>Applications:</b> | <b>WB</b> <b>IHC-P</b> <b>ELISA</b>                       |
| <b>Conjugation:</b>  | Unconjugated  |
| <b>Reactivity:</b>   | Human, Mouse, Rat   |

### Antibody Data


---

|                       |                       |
|-----------------------|-----------------------|
| <b>Gene ID:</b>       | 8773                  |
| <b>Uniprot:</b>       | AB_2760763            |
| <b>Host Species:</b>  | Rabbit                |
| <b>Purification:</b>  | Affinity purification |
| <b>Observed MW:</b>   | 23kDa                 |
| <b>Calculated MW:</b> | 23kDa                 |


## Preparation & Storage

|   |   |           |                |              |               |              |   |
|---|---|-----------|----------------|--------------|---------------|--------------|---|
| <b>Storage:</b>                           | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.<br>Store Bradford Reagent at Room Temperature for 1 Year.  |           |                |              |               |              |   |
| <b>Positive Sample:</b>                   | HeLa, Mouse kidney  |           |                |              |               |              |   |
| <b>Recommended Dilutions:</b>             | <table border="1"> <tr> <td><b>WB</b></td><td>1:500 - 1:2000</td></tr> <tr> <td><b>IHC-P</b></td><td>1:100 - 1:200</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | <b>WB</b> | 1:500 - 1:2000 | <b>IHC-P</b> | 1:100 - 1:200 | <b>ELISA</b> | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>WB</b>                                 | 1:500 - 1:2000  |           |                |              |               |              |   |
| <b>IHC-P</b>                              | 1:100 - 1:200   |           |                |              |               |              |   |
| <b>ELISA</b>                              | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.   |           |                |              |               |              |   |
| <b>Protein Quantification (Optional):</b> | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol  |           |                |              |               |              |   |


## Validation Data



Western blot analysis of various lysates using SNAP23 Rabbit pAb (CAB13909) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 2min.



Immunohistochemistry analysis of paraffin-embedded Rat lymph node using SNAP23 Rabbit pAb (CAB13909) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen using SNAP23 Rabbit pAb (CAB13909) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.