

PTCH1 Antibody

CAB14772

Description

This PTCH1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB14772

Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: PTC, BCNS, PTC1, PTCH, NBCCS, PTCH1

Clone: -

Applications:   

Conjugation: Unconjugated

Reactivity: Mouse, Rat

Antibody Data

Gene ID: 5727

Uniprot: AB_2761648

Host Species: Rabbit

Purification: Affinity purification


Observed MW: 160kDa

Calculated MW: 161kDa

Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.						
	Store Bradford Reagent at Room Temperature for 1 Year.						
Positive Sample:	Rat skeletal muscle						
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:1000</td></tr> <tr> <td>IF-P</td><td>1:50 - 1:200</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:1000	IF-P	1:50 - 1:200	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:1000						
IF-P	1:50 - 1:200						
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.						
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol						

Validation Data



Western blot analysis of lysates from Rat skeletal muscle, using PTCH1 Rabbit pAb (CAB14772) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.

Immunofluorescence analysis of paraffin-embedded mouse embryos using PTCH1 Rabbit pAb (CAB14772) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

