

## PDCD7 Monoclonal Antibody

**CAB1510**

### Description

---

This PDCD7 Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB1510

**Contents:** 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

**Category:** Monoclonal Antibody

**Synonyms:** 59K, ES18, HES18, PDCD7

**Clone:** ARC2564

**Applications:** WB IHC-P ELISA

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 10081

**Uniprot:** -

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 55kDa

**Calculated MW:** 55kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** TE-1, SK-OV-3, Mouse testis, Mouse thymus, Rat brain


**Recommended Dilutions:**

<b>WB</b>	1:500 - 1:1000
<b>IHC-P</b>	1:50 - 1:200
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data


---




Western blot analysis of various lysates using PDCD7 Rabbit mAb (CAB1510) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.




Western blot analysis of lysates from Rat brain, using PDCD7 Rabbit mAb (CAB1510) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using PDCD7 Rabbit mAb (CAB1510) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human small intestine tissue using PDCD7 Rabbit mAb (CAB1510) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using PDCD7 Rabbit mAb (CAB1510) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using PDCD7 Rabbit mAb (CAB1510) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.