

EIF5B Antibody

CAB15123

Description

This EIF5B Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB15123

Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: IF2, EIF5B

Clone: -

Applications: **WB** **IF/ICC** **ELISA**

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 9669

Uniprot: AB_2762008

Host Species: Rabbit

Purification: Affinity purification


Observed MW: 179kDa

Calculated MW: 139kDa


Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.						
	Store Bradford Reagent at Room Temperature for 1 Year.						
Positive Sample:	DU145, HeLa, Jurkat, LO2						
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>IF/ICC</td><td>1:50 - 1:200</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:2000	IF/ICC	1:50 - 1:200	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:2000						
IF/ICC	1:50 - 1:200						
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.						
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol						


Validation Data



Western blot analysis of various lysates using EIF5B Rabbit pAb (CAB15123). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 3s.



Immunofluorescence analysis of cells using EIF5B Rabbit pAb (CAB15123) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using EIF5B Rabbit pAb (CAB15123) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.