

KIF22 Antibody

CAB15285

Description

This KIF22 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB15285

Contents: 20 μ L, 100 μ L

Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: KID, OBP, KNSL4, OBP-1, OBP-2, SEMDJL2, A-328A3.2, KIF22

Clone: -

Applications: **WB** **IHC-P** **ELISA**

Conjugation: Unconjugated

Reactivity: Human, Mouse

Antibody Data

Gene ID: 3835

Uniprot: AB_2762185

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 73kDa

Calculated MW: 73kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: A431, 293T, U-87MG, A549, HeLa, Jurkat, Mouse testis

Recommended Dilutions:


WB	1:200 - 1:2000
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB15285) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 3s.



Immunohistochemistry analysis of paraffin-embedded Human colon using Rabbit pAb (CAB15285) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Rabbit pAb (CAB15285) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.