

## Beta-TRCP Antibody

**CAB1656**

### Description

---

This Beta-TRCP Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB1656

**Contents:** 20  $\mu$ L, 100  $\mu$ L

Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** FWD1, FBW1A, FBXW1, bTrCP, FBXW1A, bTrCP1, betaTrCP, BETA-TRCP,  $\beta$ -TrCP/BTRC

**Clone:** -

**Applications:** WB IHC-P IF/ICC ELISA

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 8945

**Uniprot:** AB\_2763713

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 65kDa

**Calculated MW:** 69kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** U-251MG


**Recommended Dilutions:**

<b>WB</b>	1:500 - 1:1000
<b>IHC-P</b>	1:50 - 1:200
<b>IF/ICC</b>	1:50 - 1:200
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---



Western blot analysis of lysates from U-251MG cells, using β-TrCP/BTRC Rabbit pAb (CAB1656) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (AbGn00021). Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen using  $\beta$ -TrCP/BTRC Rabbit pAb (CAB1656) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.