

## SOCS2 Antibody

CAB17035

### Description

---

This SOCS2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

|                      |                                                        |
|----------------------|--------------------------------------------------------|
| <b>SKU:</b>          | CAB17035                                               |
| <b>Contents:</b>     | 20 µL, 100 µL<br>Bradford Reagent: 1 vial (2ml)        |
| <b>Category:</b>     | Polyclonal Antibody                                    |
| <b>Synonyms:</b>     | CIS2, SSI2, Cish2, SSI-2, SOCS-2, STATI2, SOCS2        |
| <b>Clone:</b>        | -                                                      |
| <b>Applications:</b> | <span>WB</span> <span>IF/ICC</span> <span>ELISA</span> |
| <b>Conjugation:</b>  | -                                                      |
| <b>Reactivity:</b>   | Human                                                  |

### Antibody Data

---

|                       |                       |
|-----------------------|-----------------------|
| <b>Gene ID:</b>       | 8835                  |
| <b>Uniprot:</b>       | AB_2772341            |
| <b>Host Species:</b>  | Rabbit                |
| <b>Purification:</b>  | Affinity purification |
| <b>Observed MW:</b>   | 22kDa                 |
| <b>Calculated MW:</b> | 22kDa                 |

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** A-549, 293T

**Recommended Dilutions:**


|               |                                                                                                                             |
|---------------|-----------------------------------------------------------------------------------------------------------------------------|
| <b>WB</b>     | 1:500 - 1:2000                                                                                                              |
| <b>IF/ICC</b> | 1:50 - 1:100                                                                                                                |
| <b>ELISA</b>  | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data



Western blot analysis of various lysates using SOCS2 Rabbit pAb (CAB17035) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (AbGn00021). Exposure time: 90s.



Immunofluorescence analysis of U-2 OS cells using SOCS2 Rabbit pAb (CAB17035) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

