

## M1-linkage Specific Polyubiquitin Antibody

CAB18200

### Description

---

This M1-linkage Specific Polyubiquitin Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB18200

**Contents:** 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** -

**Clone:** -

**Applications:**   

**Conjugation:** -

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** -

**Uniprot:** AB\_2861977

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** Refer to figures

**Calculated MW:** -

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% sodium azide, 50% glycerol, pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** HeLa, NIH/3T3, RAW264.7, C6, Mouse brain, Mouse kidney, Rat brain, Rat kidney

**Recommended Dilutions:**

<b>WB</b>	1:500 - 1:2000
<b>DB</b>	1:500 - 1:1000
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data



Western blot analysis of various lysates using -linkage Specific Polyubiquitin Rabbit pAb (CAB18200) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.

Dot-blot analysis of all sorts of peptides using -linkage Specific Polyubiquitin antibody (CAB18200) at 1:1000 dilution.