

TBC1D4 Antibody

CAB18216

Description

This TBC1D4 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB18216
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: AS160, NIDDM5, TBC1D4
Clone: -
Applications: **WB** **IF/ICC** **ELISA**
Conjugation: -
Reactivity: Mouse, Rat


Antibody Data

Gene ID: 9882
Uniprot: AB_2861992
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 160kDa
Calculated MW: 147kDa


Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.						
	Store Bradford Reagent at Room Temperature for 1 Year.						
Positive Sample:	Rat skeletal muscle						
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>IF/ICC</td><td>1:50 - 1:200</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:2000	IF/ICC	1:50 - 1:200	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:2000						
IF/ICC	1:50 - 1:200						
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.						
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol						

Validation Data



Western blot analysis of lysates from Rat skeletal muscle, using TBC1D4 Rabbit pAb (CAB18216) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.



Immunofluorescence analysis of cells using TBC1D4 Rabbit pAb (CAB18216) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

