

## **RDH11 Antibody**

**CAB18454**

### **Description**

---

This RDH11 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### **Product Information**

---

<b>SKU:</b>	CAB18454
<b>Contents:</b>	20 µL, 100 µL Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Polyclonal Antibody
<b>Synonyms:</b>	MDT1, CGI82, PSDR1, RALR1, SCALD, ARSDR1, HCBP12, RDJCSS, SDR7C1, RDH11
<b>Clone:</b>	-
<b>Applications:</b>	<span style="background-color: red; color: white; padding: 2px 5px;">WB</span> <span style="background-color: #0070C0; color: white; padding: 2px 5px;">ELISA</span>
<b>Conjugation:</b>	-
<b>Reactivity:</b>	Human, Rat

### **Antibody Data**

---

<b>Gene ID:</b>	51109
<b>Uniprot:</b>	AB_2862221
<b>Host Species:</b>	Rabbit
<b>Purification:</b>	Affinity purification
<b>Observed MW:</b>	35kDa
<b>Calculated MW:</b>	35kDa


## Preparation & Storage

---

<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.				
	Store Bradford Reagent at Room Temperature for 1 Year.				
<b>Positive Sample:</b>	PC-3, HepG2				
<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:500 - 1:2000</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:500 - 1:2000	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:500 - 1:2000				
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.				
<b>Protein Quantification (Optional):</b>	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol				

## Validation Data

---



Western blot analysis of various lysates using Rabbit pAb (CAB18454) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.