

ZNF444 Antibody

CAB18470

Description

This ZNF444 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB18470
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: EZF2, EZF-2, ZSCAN17, ZNF444
Clone: -
Applications: **WB** | **IHC-P** | **IF/ICC** | **ELISA**
Conjugation: -
Reactivity: Human, Mouse, Rat


Antibody Data

Gene ID: 55311
Uniprot: AB_2862237
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 38kDa
Calculated MW: 35kDa

Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.								
	Store Bradford Reagent at Room Temperature for 1 Year.								
Positive Sample:	HeLa, Mouse lung, Mouse spleen								
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>IHC-P</td><td>1:50 - 1:200</td></tr> <tr> <td>IF/ICC</td><td>1:50 - 1:200</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:2000	IHC-P	1:50 - 1:200	IF/ICC	1:50 - 1:200	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:2000								
IHC-P	1:50 - 1:200								
IF/ICC	1:50 - 1:200								
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol								

Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB18470) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (AbGn00021). Exposure time: 90s.