

## WWC2 Antibody

**CAB18668**

### Description

---

This WWC2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB18668

**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** BOMB, WWC2

**Clone:** -

**Applications:** **WB** **IHC-P** **ELISA**

**Conjugation:** -

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 80014

**Uniprot:** AB\_2862405

**Host Species:** Rabbit

**Purification:** Affinity purification


**Observed MW:** 134kDa

**Calculated MW:** 134kDa


## Preparation & Storage

<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. Store Bradford Reagent at Room Temperature for 1 Year.						
<b>Positive Sample:</b>	U20S, Mouse lung						
<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:500 - 1:2000</td></tr> <tr> <td><b>IHC-P</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:500 - 1:2000	<b>IHC-P</b>	1:50 - 1:200	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:500 - 1:2000						
<b>IHC-P</b>	1:50 - 1:200						
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.						
<b>Protein Quantification (Optional):</b>	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol						

## Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB18668) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 5s.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using Rabbit pAb (CAB18668) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

