

## Fatty Acid Synthase (FASN) Monoclonal Antibody

CAB19050

### Description

---

This Fatty Acid Synthase (FASN) Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB19050

**Contents:** 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

**Category:** Monoclonal Antibody

**Synonyms:** FAS, OA-519, SDR27X1, Fatty Acid Synthase (FASN)

**Clone:** ARC0377

**Applications:**     

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 2194

**Uniprot:** AB\_2862543

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 273kDa

**Calculated MW:** 273kDa

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** A549, HeLa

**Recommended Dilutions:**


<b>WB</b>	1:1000 - 1:2000
<b>IHC-P</b>	1:500 - 1:2000
<b>IF/ICC</b>	1:100 - 1:800
<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


## Validation Data




Western blot analysis of various lysates using Fatty Acid Synthase (FASN) Rabbit mAb (CAB19050) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (AbGn00021). Exposure time: 3min.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Fatty Acid Synthase (FASN) Rabbit mAb (CAB19050) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse skin tissue using Fatty Acid Synthase (FASN) Rabbit mAb (CAB19050) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat fat tissue using Fatty Acid Synthase (FASN) Rabbit mAb (CAB19050) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Fatty Acid Synthase (FASN) Rabbit mAb (CAB19050). Western blot was performed from the immunoprecipitate using Fatty Acid Synthase (FASN) antibody (CAB19050) at a dilution of 1:500.