

TROY/TNFRSF19 Monoclonal Antibody

CAB19235

Description

This TROY/TNFRSF19 Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB19235
Contents: 20 µL, 100 µL
Bradford Reagent: 1 vial (2ml)
Category: Monoclonal Antibody
Synonyms: TAJ, TROY, TRADE, TAJ-alpha, TROY/TNFRSF19
Clone: ARC2393
Applications: WB ELISA
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat


Antibody Data

Gene ID: 55504
Uniprot: -
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 46kDa
Calculated MW: 46kDa

Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.				
Positive Sample:	22Rv1, Raji, U-251MG, Mouse lung, Mouse brain, Mouse spleen, Rat lung, Rat brain				
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:1000</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:1000	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:1000				
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.				
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol				

Validation Data



Western blot analysis of various lysates using TROY/TNFRSF19 Rabbit mAb (CAB19235) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 1s.