

CIRBP Monoclonal Antibody

CAB19290

Description

This CIRBP Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB19290
Contents: 20 µL, 100 µL
Bradford Reagent: 1 vial (2ml)
Category: Monoclonal Antibody
Synonyms: CIRP, CIRBP
Clone: ARC2472
Applications: **WB** | **IHC-P** | **IP** | **ELISA** | **IF-P**
Conjugation: Unconjugated
Reactivity: Human

Antibody Data

Gene ID: 1153
Uniprot: -
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 19kDa
Calculated MW: 19kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: 293T, HepG2, SH-SY5Y

Recommended Dilutions:


WB	1:500 - 1:2000
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cell
IF-P	1:50 - 1:200
IHC-P	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data




Western blot analysis of various lysates using CIRBP Rabbit mAb (CAB19290) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human pancreas tissue using CIRBP Rabbit mAb (CAB19290) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human thyroid tissue using CIRBP Rabbit mAb (CAB19290) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Human breast cancer tissue using CIRBP Rabbit mAb (CAB19290, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (CABS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.