Musashi-2 (MSI2) Monoclonal Antibody
CAB19814

Description

This Musashi-2 (MSI2) Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	CAB19814
Contents:	20 µL, 100 µL
	Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Synonyms:	MSI2H, Musashi-2 (MSI2)
Clone:	ARC2341
Applications:	WB IHC-P ELISA
Conjugation:	Unconjugated
Reactivity:	Human, Mouse, Rat

Antibody Data

Gene ID:	124540
Uniprot:	-
Host Species:	Rabbit
Purification:	Affinity purification
Observed MW:	35kDa
Calculated MW:	35kDa

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Preparation & Storage

|	
Storage:	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3. Store Bradford Reagent at Room Temperature for 1 Year.
Positive Sample:	HeLa, HT-29, PC-12, MCF7, Mouse brain, Rat brain

Recommended Dilutions:

WB	1:1000 - 1:6000
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol

Validation Data

Western blot analysis of various lysates using Musashi-2 Rabbit mAb (CAB19814) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 5s.

Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Musashi-2 Rabbit mAb (CAB19814) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.