

## Ajuba Polyclonal Antibody

CAB21760

### Description

---

This Ajuba Polyclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB21760  
**Contents:** 20 µL, 100 µL  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** JUB, ba  
**Clone:** -  
**Applications:** WB ELISA  
**Conjugation:** Unconjugated  
**Reactivity:** Human, Rat

### Antibody Data

---

**Gene ID:** 84962  
**Uniprot:** -  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 55kDa/  
**Calculated MW:** 57kDa

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** HepG2, Rat kidney, Rat kidney , Hep G2

**Recommended Dilutions:**


WB	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data



Western blot analysis of lysates from HepG2 cells, using [KO Validated] Ajuba Rabbit pAb (CAB21760) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.



Western blot analysis of lysates from Rat kidney using [KO Validated] Ajuba Rabbit pAb (CAB21760) at 1:1000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 180s.

