

TriMethyl-Histone H3-K27 Antibody

CAB2363

Description

This TriMethyl-Histone H3-K27 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB2363

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: H3t, H3.4, H3/g, H3FT, H3C16, HIST3H3, TriMethyl-Histone H3-K27

Clone: -

Applications: WB IHC-P IF/ICC IP ChIP ChIP-seq ELISA

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat, Other (Wide Range Predicted)

Antibody Data

Gene ID: 8290 8350

Uniprot: AB_2756439

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 17 kDa

Calculated MW: 15 kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: MCF7 acid extract, 293T acid extract, C2C12 acid extract, NIH/3T3 acid extract, C6 acid extract


Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:5000</td></tr> <tr> <td>IP</td><td>0.5ug-4ug antibody for 400ug-600ug extracts of whole cells</td></tr> <tr> <td>IF/ICC</td><td>1:50 - 1:2000</td></tr> <tr> <td>IHC-P</td><td>1:50 - 1:200</td></tr> <tr> <td>DB</td><td>1:500 - 1:100 ChIP 5µg antibody for 5µg-10µg of Chromatin ChIP-seq 1:20 - 1:100</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:5000	IP	0.5ug-4ug antibody for 400ug-600ug extracts of whole cells	IF/ICC	1:50 - 1:2000	IHC-P	1:50 - 1:200	DB	1:500 - 1:100 ChIP 5µg antibody for 5µg-10µg of Chromatin ChIP-seq 1:20 - 1:100	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:5000												
IP	0.5ug-4ug antibody for 400ug-600ug extracts of whole cells												
IF/ICC	1:50 - 1:2000												
IHC-P	1:50 - 1:200												
DB	1:500 - 1:100 ChIP 5µg antibody for 5µg-10µg of Chromatin ChIP-seq 1:20 - 1:100												
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.												

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data




Western blot analysis of various lysates using TriMethyl-Histone - Rabbit pAb (CAB2363) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 0.5s.



Dot-blot analysis of all sorts of methylation peptides using TriMethyl-Histone - antibody (CAB2363).



Immunofluorescence analysis of PC-12 cells using TriMethyl-Histone - Rabbit pAb (CAB2363) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TriMethyl-Histone - antibody (CAB2363) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.