

MonoMethyl-Histone H3-R17 Antibody

CAB3151

Description

This MonoMethyl-Histone H3-R17 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

| | |
|----------------------|---|
| SKU: | CAB3151 |
| Contents: | 20 µL, 100 µL Bradford Reagent: 1 vial (2ml) |
| Category: | Polyclonal Antibody |
| Synonyms: | H3/j, H3C1, H3C2, H3C3, H3C4, H3C6, H3C7, H3C8, H3FJ, H3C10, H3C11, HIST1H3J, MonoMethyl-Histone H3-R17 |
| Clone: | - |
| Applications: | WB IHC-P IF/ICC ELISA |
| Conjugation: | Unconjugated |
| Reactivity: | Human, Mouse, Rat, Other (Wide Range Predicted) |

Antibody Data

| | |
|-----------------------|-----------------------|
| Gene ID: | 8290 8350 |
| Uniprot: | AB_2764945 |
| Host Species: | Rabbit |
| Purification: | Affinity purification |
| Observed MW: | 17kDa |
| Calculated MW: | 15kDa |

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: C6


| | | |
|-------------------------------|---------------|---|
| Recommended Dilutions: | WB | 1:500 - 1:1000 |
| | IHC-P | 1:50 - 1:200 |
| | IF/ICC | 1:50 - 1:200 |
| | ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data




Western blot analysis of lysates from cells, using MonoMethyl-Histone - Rabbit pAb (CAB3151) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (AbGn00021). Exposure time: 180s.




Dot-blot analysis of all sorts of methylation peptides using MonoMethyl-Histone - antibody (CAB3151).




Immunohistochemistry analysis of paraffin-embedded Human breast using MonoMethyl-Histone - Rabbit pAb (CAB3151) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using MonoMethyl-Histone - Rabbit pAb (CAB3151) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of NIH-3T3 cells using MonoMethyl-Histone - Rabbit pAb (CAB3151) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using MonoMethyl-Histone - Rabbit pAb (CAB3151) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.