

CLPP Monoclonal Antibody

CAB3214

Description

This CLPP Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB3214

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Monoclonal Antibody

Synonyms: DFNB81, PRLTS3, CLPP

Clone: ARC1929

Applications: WB IHC-P IF/ICC ELISA

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 8192

Uniprot: AB_2863027

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 26kDa

Calculated MW: 30kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: HeLa, SW620, Hep G2, Mouse liver, Mouse heart, Rat liver


Recommended Dilutions:	WB	1:500 - 1:1000
	IHC-P	1:50 - 1:200
	IF/ICC	1:50 - 1:200
	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data



Western blot analysis of various lysates using CLPP Rabbit mAb (CAB3214) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 3min.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using CLPP Rabbit mAb (CAB3214) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunofluorescence analysis of NIH/3T3 cells using CLPP Rabbit mAb (CAB3214) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.