

## DOCK2 Antibody

**CAB3595**

### Description

---

This DOCK2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB3595  
**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** IMD40, DOCK2  
**Clone:** -  
**Applications:** **WB** **IF/ICC** **IP** **ELISA**  
**Conjugation:** Unconjugated  
**Reactivity:** Human, Mouse

### Antibody Data

---

**Gene ID:** 1794  
**Uniprot:** AB\_2765171  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 240kDa  
**Calculated MW:** 212kDa


## Preparation & Storage

---


<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. Store Bradford Reagent at Room Temperature for 1 Year.								
<b>Positive Sample:</b>	Jurkat								
<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:500 - 1:1000</td></tr> <tr> <td><b>IF/ICC</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>IP</b></td><td>0.5µg-4µg antibody for 200µg-400µg extracts of whole cells</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:500 - 1:1000	<b>IF/ICC</b>	1:50 - 1:200	<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:500 - 1:1000								
<b>IF/ICC</b>	1:50 - 1:200								
<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells								
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								
<b>Protein Quantification (Optional):</b>	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol								

## Validation Data

---



Western blot analysis of lysates from Jurkat cells, using DOCK2 Rabbit pAb (CAB3595). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of cells using DOCK2 Rabbit pAb (CAB3595) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.