

## TCP1 alpha Monoclonal Antibody

**CAB3887**

### Description

---

This TCP1 alpha Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB3887

**Contents:** 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

**Category:** Monoclonal Antibody

**Synonyms:** CCT1, CCTa, D6S230E, CCT-alpha, TCP-1-alpha, TCP1 alpha

**Clone:** ARC0859

**Applications:** WB IHC-P ELISA

**Conjugation:** Unconjugated

**Reactivity:** Human

### Antibody Data

---

**Gene ID:** 6950

**Uniprot:** -

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 60kDa

**Calculated MW:** 60kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** Jurkat


**Recommended Dilutions:**

<b>WB</b>	1:500 - 1:1000
<b>IHC-P</b>	1:50 - 1:200
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---



Western blot analysis of lysates from Jurkat cells, using alpha Rabbit mAb (CAB3887) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human colon using alpha Rabbit mAb (CAB3887) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.