

Lipoprotein lipase (LPL) Monoclonal Antibody

CAB4115

Description

This Lipoprotein lipase (LPL) Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB4115

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Monoclonal Antibody

Synonyms: LIPD, HDLCQ11, Lipoprotein lipase (LPL)

Clone: ARC0904

Applications:  

Conjugation: Unconjugated

Reactivity: Human, Mouse

Antibody Data

Gene ID: 4023

Uniprot: AB_2863190

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 53kDa

Calculated MW: 53kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: Human plasma, Mouse liver

Recommended Dilutions:


WB	1:1000 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Western blot analysis of lysates from Human plasma, using Lipoprotein lipase (LPL) (LPL) Rabbit mAb (CAB4115) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 3s.



Western blot analysis of lysates from Mouse liver, using Lipoprotein lipase (LPL) (LPL) Rabbit mAb (CAB4115) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 3min.

