

## VARS Antibody

**CAB4182**

### Description

---

This VARS Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB4182  
**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** G7A, VARS, VARS2, NDMSCA  
**Clone:** -  
**Applications:** **WB** | **IHC-P** | **IF/ICC** | **ELISA**  
**Conjugation:** Unconjugated  
**Reactivity:** Human, Mouse, Rat

### Antibody Data


---

**Gene ID:** 7407  
**Uniprot:** AB\_2765544  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 140kDa  
**Calculated MW:** 140kDa


## Preparation & Storage

<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.								
	Store Bradford Reagent at Room Temperature for 1 Year.								
<b>Positive Sample:</b>	HepG2, HL-60, K-562, SW620, Mouse brain, Mouse heart, Mouse liver, Rat liver								
<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:500 - 1:2000</td></tr> <tr> <td><b>IHC-P</b></td><td>1:50 - 1:100</td></tr> <tr> <td><b>IF/ICC</b></td><td>1:50 - 1:100</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:500 - 1:2000	<b>IHC-P</b>	1:50 - 1:100	<b>IF/ICC</b>	1:50 - 1:100	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:500 - 1:2000								
<b>IHC-P</b>	1:50 - 1:100								
<b>IF/ICC</b>	1:50 - 1:100								
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								
<b>Protein Quantification (Optional):</b>	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol								


## Validation Data




Western blot analysis of various lysates using VARS Rabbit pAb (CAB4182) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using VARS Rabbit pAb (CAB4182) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using VARS Rabbit pAb (CAB4182) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of cells using VARS Rabbit pAb (CAB4182) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.