

## Cytokeratin 7 (KRT7) Monoclonal Antibody

CAB4765

### Description

---

This Cytokeratin 7 (KRT7) Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB4765

**Contents:** 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

**Category:** Monoclonal Antibody

**Synonyms:** K7, CK7, SCL, K2C7, Cytokeratin 7 (KRT7)

**Clone:** ARC1267

**Applications:** WB IHC-P ELISA IF-P

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 3855

**Uniprot:** AB\_2863343

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 55kDa

**Calculated MW:** 51kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** HeLa, A-549, Mouse lung, Mouse kidney, Rat kidney

<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:1000 - 1:4000</td></tr> <tr> <td><b>IF-P</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>IHC-P</b></td><td>1:500 - 1:2000</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:1000 - 1:4000	<b>IF-P</b>	1:50 - 1:200	<b>IHC-P</b>	1:500 - 1:2000	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:1000 - 1:4000								
<b>IF-P</b>	1:50 - 1:200								
<b>IHC-P</b>	1:500 - 1:2000								
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data


---




Western blot analysis of various lysates using Cytokeratin 7 Rabbit mAb (CAB4765) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.




Western blot analysis of various lysates using Cytokeratin 7 Rabbit mAb (CAB4765) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 3min.




Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using Cytokeratin 7 Rabbit mAb (CAB4765) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer tissue using Cytokeratin 7 Rabbit mAb (CAB4765) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta tissue using Cytokeratin 7 Rabbit mAb (CAB4765) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Cytokeratin 7 Rabbit mAb (CAB4765) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.