

P23/PTGES3 Antibody

CAB5325

Description

This P23/PTGES3 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

| | |
|----------------------|---|
| SKU: | CAB5325 |
| Contents: | 20 µL, 100 µL Bradford Reagent: 1 vial (2ml) |
| Category: | Polyclonal Antibody |
| Synonyms: | P23, TEBP, cPGES, P23/PTGES3 |
| Clone: | - |
| Applications: | WB ELISA |
| Conjugation: | Unconjugated |
| Reactivity: | Human, Mouse, Rat |


Antibody Data

| | |
|-----------------------|-----------------------|
| Gene ID: | 10728 |
| Uniprot: | AB_2766137 |
| Host Species: | Rabbit |
| Purification: | Affinity purification |
| Observed MW: | 23kDa |
| Calculated MW: | 19kDa |

Preparation & Storage

| | | | | | |
|---|--|-----------|----------------|--------------|---|
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. | | | | |
| | Store Bradford Reagent at Room Temperature for 1 Year. | | | | |
| Positive Sample: | HeLa, Jurkat, Mouse testis, Rat testis | | | | |
| Recommended Dilutions: | <table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | WB | 1:500 - 1:2000 | ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| WB | 1:500 - 1:2000 | | | | |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. | | | | |
| Protein Quantification (Optional): | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol | | | | |

Validation Data



Western blot analysis of various lysates using /PTGES3 Rabbit pAb (CAB5325) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (AbGn00021). Exposure time: 90s.