

NOGO Receptor 1 (RTN4R) Antibody

CAB5847

Description

This NOGO Receptor 1 (RTN4R) Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB5847
Contents: 20 µL, 100 µL
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: NGR, NOGOR, NOGO Receptor 1 (RTN4R)
Clone: -
Applications: WB IF/ICC ELISA
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 65078
Uniprot: AB_2766597
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 66kDa
Calculated MW: 51kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: C6, Rat uterus

Recommended Dilutions:


WB	1:100 - 1:500
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Western blot analysis of various lysates using NOGO Receptor 1 (RTN4R) Rabbit pAb (CAB5847) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 180s.



Immunofluorescence analysis of Neuro-2a cells using NOGO Receptor 1 (RTN4R) Rabbit pAb (CAB5847) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.