

## TNFRSF10D Antibody

**CAB6136**

### Description

---

This TNFRSF10D Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB6136

**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** DCR2, CD264, TRUNDD, TRAILR4, TRAIL-R4, TNFRSF10D

**Clone:** -

**Applications:** **WB** **IHC-P** **ELISA**

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse

### Antibody Data

---

**Gene ID:** 8793

**Uniprot:** AB\_2766766

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 47kDa

**Calculated MW:** 42kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** THP-1


**Recommended Dilutions:**

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---



Western blot analysis of lysates from THP-1 cells, using TNFRSF10D Rabbit pAb (CAB6136) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using TNFRSF10D Rabbit pAb (CAB6136) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.