

## PPIE Antibody

CAB7061

### Description

---

This PPIE Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB7061  
**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** CYP33, CYP-33, PPIE  
**Clone:** -  
**Applications:** WB IP ELISA  
**Conjugation:** Unconjugated  
**Reactivity:** Human, Mouse

### Antibody Data

---

**Gene ID:** 10450  
**Uniprot:** AB\_2767616  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 33kDa  
**Calculated MW:** 33kDa

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** HeLa, THP-1, Jurkat, HL-60, Mouse testis, Mouse thymus


<b>Recommended Dilutions:</b>	<b>WB</b>	1:500 - 1:2000
	<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data



Western blot analysis of various lysates using PPIE Rabbit pAb (CAB7061) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.



Immunoprecipitation analysis of 200 µg extracts of HeLa cells using 1 µg PPIE antibody (CAB7061). Western blot was performed from the immunoprecipitate using PPIE antibody (CAB7061) at a dilution of 1:1000.