

PHIP Antibody

CAB7207

Description

This PHIP Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB7207

Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: ndrp, BRWD2, DIDOD, WDR11, DCAF14, CHUJANS, PHIP

Clone: -

Applications: **WB** **IHC-P** **IF/ICC** **ELISA**

Conjugation: Unconjugated

Reactivity: Human, Mouse

Antibody Data

Gene ID: 55023

Uniprot: AB_2767756

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 250kDa

Calculated MW: 207kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: HeLa, Jurkat

Recommended Dilutions:


| | |
|---------------|---|
| WB | 1:500 - 1:2000 |
| IHC-P | 1:50 - 1:200 |
| IF/ICC | 1:50 - 1:100 |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data




Western blot analysis of various lysates using PHIP Rabbit pAb (CAB7207) at 1:1000 dilution._Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution._Lysates/proteins: 25µg per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit (AbGn00021)._Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human well-differentiated squamous skin carcinoma using PHIP Rabbit pAb (CAB7207) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse tumor using PHIP Rabbit pAb (CAB7207) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of MCF-7 cells using PHIP Rabbit pAb (CAB7207). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution.