

## Acetyl-Histone H3-K56 Antibody

**CAB7256**

### Description

---

This Acetyl-Histone H3-K56 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB7256

**Contents:** 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** H3t, H3.4, H3/g, H3FT, H3C16, HIST3H3, Acetyl-Histone H3-K56

**Clone:** -

**Applications:** WB IF/ICC ChIP ELISA

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat, Other (Wide Range Predicted)

### Antibody Data

---

**Gene ID:** 8290 8350

**Uniprot:** AB\_2767800

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 17 kDa

**Calculated MW:** 15 kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** HeLa treated with TSA, NIH/3T3 treated with TSA, C6 treated with TSA

<b>Recommended Dilutions:</b>	<b>WB</b>	1:1000 - 1:10000
	<b>IF/ICC</b>	1:50 - 1:200
	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. ChIP 5µg antibody for 5µg-10µg of Chromatin


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data


---




Western blot analysis of lysates from HeLa cells, using Acetyl-Histone - Rabbit pAb (CAB7256) at 1:10000 dilution. HeLa cells were treated with TSA (1 uM) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.




Western blot analysis of lysates from NIH/3T3 cells, using Acetyl-Histone - Rabbit pAb (CAB7256) at 1:10000 dilution. NIH/3T3 cells were treated with TSA (1  $\mu$ M) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.




Immunofluorescence analysis of treated with TSA jia cells using Acetyl-Histone - Rabbit pAb (CAB7256) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of treated with TSA jia cells using Acetyl-Histone - Rabbit pAb (CAB7256) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS treated with TSA jia U-2 OS cells using Acetyl-Histone - Rabbit pAb (CAB7256) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Chromatin immunoprecipitation analysis of extracts of Hale cells, using Acetyl-Histone - antibody (CAB7256) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.