

RNF7 Antibody

CAB7301

Description

This RNF7 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB7301
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: SAG, ROC2, rbx2, CKBBP1, RNF7
Clone: -
Applications: **WB** **IF/ICC** **ELISA**
Conjugation: Unconjugated
Reactivity: Human

Antibody Data

Gene ID: 9616
Uniprot: AB_2767842
Host Species: Rabbit
Purification: Affinity purification
Observed MW: Refer to figures
Calculated MW: 13kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample:

Recommended Dilutions:	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px;">WB</td><td style="padding: 5px;">1:500 - 1:2000</td></tr> <tr> <td style="padding: 5px;">IF/ICC</td><td style="padding: 5px;">1:20 - 1:50</td></tr> <tr> <td style="padding: 5px;">ELISA</td><td style="padding: 5px;">Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:2000	IF/ICC	1:20 - 1:50	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:2000						
IF/ICC	1:20 - 1:50						
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.						

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Immunofluorescence analysis of MCF-7 cells using Rabbit pAb (CAB7301). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.