

## hnRNP A1 Antibody

**CAB7491**

### Description

---

This hnRNP A1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB7491

**Contents:** 20  $\mu$ L, 100  $\mu$ L

Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** UP 1, ALS19, ALS20, HNRPA1, IBMPFD3, HNRPA1L3, hnRNP A1, hnRNP-A1

**Clone:** -

**Applications:** **WB** **IHC-P** **IF/ICC** **ELISA**

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 3178

**Uniprot:** AB\_2768022

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 39kDa

**Calculated MW:** 39kDa

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** A-549, Jurkat, HepG2, HL-60, 293T


<b>Recommended Dilutions:</b>	<b>WB</b>	1:500 - 1:2000
	<b>IHC-P</b>	1:50 - 1:200
	<b>IF/ICC</b>	1:50 - 1:200
	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


## Validation Data




Western blot analysis of various lysates using hnRNP Rabbit pAb (CAB7491) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.




Western blot analysis of various lysates using hnRNP Rabbit pAb (CAB7491) at 1:200 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 5min.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using hnRNP Rabbit pAb (CAB7491) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using hnRNP Rabbit pAb (CAB7491) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of cells using hnRNP Rabbit pAb (CAB7491). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution.