

## PRKDC Antibody

CAB7716

### Description

---

This PRKDC Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB7716

**Contents:** 20  $\mu$ L, 100  $\mu$ L

Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** HYRC, p350, DNAPK, DNPK1, HYRC1, IMD26, XRCC7, DNAPKc, DNA-PK, DNA-PKcs

**Clone:** -

**Applications:** WB IHC-P IF/ICC ELISA

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse

### Antibody Data

---

**Gene ID:** 5591

**Uniprot:** AB\_2771830

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 469kDa

**Calculated MW:** 469kDa

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** HepG2, HeLa, A-549

**Recommended Dilutions:**


<b>WB</b>	1:500 - 1:2000
<b>IHC-P</b>	1:50 - 1:200
<b>IF/ICC</b>	1:50 - 1:200
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


## Validation Data




Western blot analysis of various lysates using DNA-PKcs Rabbit pAb (CAB7716) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human well-differentiated squamous skin carcinoma using DNA-PKcs Rabbit pAb (CAB7716) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver using DNA-PKcs Rabbit pAb (CAB7716) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of U-2OS cells using DNA-PKcs Rabbit pAb (CAB7716) at dilution of 1:100. Blue: DAPI for nuclear staining.