

ECSIT Antibody

CAB7804

Description

This ECSIT Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB7804
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: SITPEC, ECSIT
Clone: -
Applications: **WB** **IF/ICC** **ELISA**
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 51295
Uniprot: AB_2769260
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 49kDa
Calculated MW: 49kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: HepG2

Recommended Dilutions:


WB	1:500 - 1:2000
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data



Western blot analysis of lysates from HepG2 cells, using ECSIT Rabbit pAb (CAB7804) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.



Immunofluorescence analysis of H9C2 cells using ECSIT Rabbit pAb (CAB7804) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using ECSIT Rabbit pAb (CAB7804) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.