

## P2RY14 Antibody

**CAB8178**

### Description

---

This P2RY14 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB8178

**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** P2Y14, BPR105, GPR105, P2RY14

**Clone:** -

**Applications:** **WB** **IHC-P** **ELISA**

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 9934

**Uniprot:** AB\_2770762

**Host Species:** Rabbit

**Purification:** Affinity purification

**Observed MW:** 38kDa

**Calculated MW:** 39kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** K-562


**Recommended Dilutions:**

<b>WB</b>	1:500 - 1:2000
<b>IHC-P</b>	1:50 - 1:200
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---



Western blot analysis of lysates from K-562 cells, using P2RY14 Rabbit pAb (CAB8178) at 1:1500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Rat ovary using P2RY14 Rabbit pAb (CAB8178) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.