

## EIF3J Antibody

**CAB8637**

### Description

---

This EIF3J Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB8637

**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)

**Category:** Polyclonal Antibody

**Synonyms:** EIF3S1, eIF3-p35, eIF3-alpha, EIF3J

**Clone:** -

**Applications:** **WB** **IF/ICC** **ELISA**

**Conjugation:** Unconjugated

**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 8669

**Uniprot:** AB\_2769292

**Host Species:** Rabbit

**Purification:** Affinity purification


**Observed MW:** 37kDa

**Calculated MW:** 29kDa


## Preparation & Storage

|   |   |           |                |               |              |              |   |
|---|---|-----------|----------------|---------------|--------------|--------------|---|
| <b>Storage:</b>                           | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.  |           |                |               |              |              |   |
|   | Store Bradford Reagent at Room Temperature for 1 Year.  |           |                |               |              |              |   |
| <b>Positive Sample:</b>                   | SKOV3, HeLa, Jurkat, A-549, U-87MG, LO2, Mouse testis, Mouse liver  |           |                |               |              |              |   |
| <b>Recommended Dilutions:</b>             | <table border="1"> <tr> <td><b>WB</b></td><td>1:200 - 1:2000</td></tr> <tr> <td><b>IF/ICC</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | <b>WB</b> | 1:200 - 1:2000 | <b>IF/ICC</b> | 1:50 - 1:200 | <b>ELISA</b> | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>WB</b>                                 | 1:200 - 1:2000  |           |                |               |              |              |   |
| <b>IF/ICC</b>                             | 1:50 - 1:200  |           |                |               |              |              |   |
| <b>ELISA</b>                              | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.   |           |                |               |              |              |   |
| <b>Protein Quantification (Optional):</b> | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol  |           |                |               |              |              |   |


## Validation Data



Western blot analysis of various lysates using EIF3J Rabbit pAb (CAB8637) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.



Immunofluorescence analysis of HeLa cells using EIF3J Rabbit pAb (CAB8637) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using EIF3J Rabbit pAb (CAB8637) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.