

elf3B Monoclonal Antibody

CAB9143

Description

This elf3B Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB9143

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Monoclonal Antibody

Synonyms: PRT1, EIF3S9, EIF3-ETA, EIF3-P110, EIF3-P116, elf3B

Clone: ARC1448

Applications: WB IF/ICC ELISA

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 8662

Uniprot: AB_2863668

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 116kDa

Calculated MW: 93kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: HeLa, Jurkat

Recommended Dilutions:


| | |
|--------|---|
| WB | 1:500 - 1:1000 |
| IF/ICC | 1:50 - 1:200 |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Western blot analysis of various lysates using eIF3B Rabbit mAb (CAB9143) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 1s.



Confocal imaging of NIH/3T3 cells using eIF3B Rabbit mAb (CAB9143, dilution 1:100) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (CABS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (CABC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.