

EYA1 Antibody

CAB9534

Description

This EYA1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB9534
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: BOP, BOR, BOS1, OFC1, EYA1
Clone: -
Applications: **WB** **IHC-P** **ELISA**
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat


Antibody Data

Gene ID: 2138
Uniprot: AB_2769383
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 55kDa
Calculated MW: 65kDa


Preparation & Storage

| | | | | | | | |
|---|--|-----------|----------------|--------------|--------------|--------------|---|
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3. | | | | | | |
| | Store Bradford Reagent at Room Temperature for 1 Year. | | | | | | |
| Positive Sample: | SH-SY5Y, 293T, Mouse testis, Mouse brain, Rat brain | | | | | | |
| Recommended Dilutions: | <table border="1"> <tr> <td>WB</td><td>1:200 - 1:2000</td></tr> <tr> <td>IHC-P</td><td>1:50 - 1:200</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | WB | 1:200 - 1:2000 | IHC-P | 1:50 - 1:200 | ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| WB | 1:200 - 1:2000 | | | | | | |
| IHC-P | 1:50 - 1:200 | | | | | | |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. | | | | | | |
| Protein Quantification (Optional): | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol | | | | | | |

Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB9534) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human placenta using Rabbit pAb (CAB9534) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

