

Torsin A Monoclonal Antibody

CAB9579

Description

This Torsin A Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB9579
Contents: 20 µL, 100 µL
Bradford Reagent: 1 vial (2ml)
Category: Monoclonal Antibody
Synonyms: DQ2, AMC5, DYT1, TorsinA/TOR1A
Clone: ARC1645
Applications: WB IF/ICC ELISA
Conjugation: Unconjugated
Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 1861
Uniprot: -
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 38kDa
Calculated MW: 38kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: 293T, Hep G2, U-87 MG, Mouse liver, Rat liver


Recommended Dilutions:	WB	1:500 - 1:1000
	IF/ICC	1:100 - 1:400
	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data



Western blot analysis of various lysates using TorsinA/TOR1A Rabbit mAb (CAB9579) at 1:500 dilution incubated at room temperature for 1.5 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 60s.



Confocal imaging of cells using TorsinA/TOR1A Rabbit mAb (CAB9579, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (CABS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (CABC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.