

## Mouse Control IgG

CABC011


### Description

---

This Mouse Control IgG is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

|                      |   |
|----------------------|---|
| <b>SKU:</b>          | CABC011   |
| <b>Contents:</b>     | 20 $\mu$ L, 100 $\mu$ L<br>Bradford Reagent: 1 vial (2ml)                           |
| <b>Category:</b>     |   |
| <b>Synonyms:</b>     | -   |
| <b>Clone:</b>        | -   |
| <b>Applications:</b> |  |
| <b>Conjugation:</b>  | Unconjugated  |
| <b>Reactivity:</b>   | -   |

### Antibody Data

---

|                       |                         |
|-----------------------|-------------------------|
| <b>Gene ID:</b>       | -                       |
| <b>Uniprot:</b>       | AB_2770414              |
| <b>Host Species:</b>  | Mouse                   |
| <b>Purification:</b>  | Potein A/G purification |
| <b>Observed MW:</b>   | -                       |
| <b>Calculated MW:</b> | -                       |


## Preparation & Storage

---

|   |  |
|---|--|
| <b>Storage:</b>                           | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH 7.3.<br>Store Bradford Reagent at Room Temperature for 1 Year.   |
| <b>Positive Sample:</b>                   | 293T   |
| <b>Recommended Dilutions:</b>             | N/A  |
| <b>Protein Quantification (Optional):</b> | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol |

## Validation Data

---



Immunoprecipitation of over-expressed Myc-tagged protein in 293T cells incubated using Myc-tag antibody (CABE010). A mock served as negative control using mouse Control IgG (CABC011) and over-expressed 293T cell lysate served as positive control.