

Phospho-POLR2A-S5 Monoclonal Antibody

CABP0997

Description

This Phospho-POLR2A-S5 Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	CABP0997
Contents:	20 µL, 100 µL Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Synonyms:	RPB1, RPO2, POLR2, POLRA, RPBl1, RPOL2, NEDHIB, RplIILS, hsRPB1, hRPB220, Phospho-POLR2A CTD-S5
Clone:	ARC1541
Applications:	WB IHC-P IF/ICC ELISA
Conjugation:	Unconjugated
Reactivity:	Human, Mouse, Rat

Antibody Data

Gene ID:	5430
Uniprot:	AB_2863889
Host Species:	Rabbit
Purification:	Affinity purification
Observed MW:	270kDa
Calculated MW:	217kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


Positive Sample: MCF7 treated with Doxorubicin, MCF7 treated with nocodazole, C2C12

Recommended Dilutions:


WB	1:1000 - 1:4000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data




Western blot analysis of various lysates using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at 1:1000 dilution. The blot shows bands for p-POLR2A CTD-S55 and β-actin across different cell lines and treatments. Lanes are grouped by cell line (MCF7 and C2C12) and treatment (Doxorubicin, Nocodazole, CIP). The treatments are indicated by '+' or '-' below the lanes.




Immunohistochemistry analysis of paraffin-embedded Mouse lung tissue using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Phospho-POLR2A CTD-Rabbit mAb (CABP0997) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using Phospho-POLR2A CTD-Rabbit mAb (CABP0997) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using Phospho-POLR2A CTD-Rabbit mAb (CABP0997) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of A-549 cells using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Phospho-POLR2A CTD- Rabbit mAb (CABP0997) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.