

## CARD17 Antibody

PACO19390

### Description

---

This CARD17 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** PACO19390

**Contents:** 50µl

Bradford Reagent: 1 vial (2ml)

**Category:** -

**Synonyms:** CAR17\_HUMAN antibody, CARD 17 antibody, CARD17 antibody, Caspase 1 inhibitor antibody, Caspase recruitment domain containing protein 17 antibody, Caspase recruitment domain-containing protein 17 antibody, Caspase-1 inhibitor INCA antibody, Caspase1 inhibitor antibody, Inhibitory caspase recruitment domain (CARD) protein antibody, Inhibitory caspase recruitment domain protein antibody

**Clone:** -

**Applications:** **ELISA** **IHC**

**Conjugation:** Non-conjugated

**Reactivity:** Human

### Antibody Data

---

**Isotype:** IgG

**Uniprot:** Q5XLA6

**Host Species:** Rabbit

**Purification:** Antigen affinity purification

**Immunogen:** Synthetic peptide of Human CARD17

**Immunogen Species:** Homo sapiens (Human)

**Buffer:** -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Form:** Liquid

## Preparation & Storage

---

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.


<b>Recommended Dilutions:</b>	<b>Application</b>	<b>Recommended Dilution</b>
	ELISA	1:1000-1:5000
	IHC	1:50-1:200

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---

### Image



### Description

The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO19390 at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification:  $\times 200$ )

The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO19390 at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification:  $\times 200$ )