

DYRK3 Antibody

PACO26365

Description

This DYRK3 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: PACO26365

Contents: 50µg
Bradford Reagent: 1 vial (2ml)

Category: -

Synonyms: Dual specificity tyrosine (Y) phosphorylation regulated kinase 3 antibody, Dual specificity tyrosine phosphorylation-regulated kinase 3 antibody, Dual specificity tyrosine-phosphorylation-regulated kinase 3 antibody, Dyrk3 antibody, DYRK3_HUMAN antibody, DYRK5 antibody, hYAK3 2 antibody, Protein kinase Dyrk3 antibody, RED antibody, REDK antibody, Regulatory erythroid kinase antibody

Clone: Polyclonal

Applications: **ELISA** **IHC**

Conjugation: Non-conjugated

Reactivity: Human

Antibody Data

Isotype: IgG

Uniprot: O43781

Host Species: Rabbit

Purification: Antigen Affinity Purified

Immunogen: Recombinant Human Dual specificity tyrosine-phosphorylation-regulated kinase 3 protein (1-300AA)

Immunogen Species: Homo sapiens (Human)

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:20-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

IHC image of PACO26365 diluted at 1:100 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.