

KYAT3 Antibody

PACO36714

Description

This KYAT3 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: PACO36714

Contents: 50µg

Bradford Reagent: 1 vial (2ml)

Category: -

Synonyms: CCBL 2 antibody, Cysteine conjugate beta lyase 2 antibody, Cysteine S conjugate beta lyase 2 antibody, Cysteine-S-conjugate beta-lyase 2 antibody, DKFZp547N1117 antibody, DKFZp667D0223 antibody, KAT 3 antibody, KAT III antibody, KAT3 antibody, KAT3_HUMAN antibody, KATIII antibody, kyat3 antibody, Kynurenine aminotransferase 3 antibody, Kynurenine aminotransferase III antibody, Kynurenine oxoglutarate transaminase 3 antibody, Kynurenine oxoglutarate transaminase III antibody, Kynurenine--glyoxylate transaminase antibody, Kynurenine--oxoglutarate transaminase 3 antibody, Kynurenine--oxoglutarate...Read more

Clone: Polyclonal

Applications: **ELISA** **IHC**

Conjugation: Non-conjugated

Reactivity: Human

Antibody Data

Isotype: IgG

Uniprot: Q6YP21

Host Species: Rabbit

Purification: >95%, Protein G purified

Immunogen: Recombinant Human Kynurenine--oxoglutarate transaminase 3 protein (1-454AA)

Immunogen Species: Homo sapiens (Human)

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Form: Liquid

Preparation & Storage

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Store Bradford Reagent at Room Temperature for 1 Year.


Recommended Dilutions:

Application	Recommended Dilution
IHC	1:20-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

Immunohistochemistry of paraffin-embedded human adrenal gland tissue using PACO36714 at dilution of 1:100

Immunohistochemistry of paraffin-embedded human small intestine tissue using PACO36714 at dilution of 1:100