

ARPIN Antibody

PACO37454

Description

This ARPIN Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO37454
Contents:	50µg Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	2610034B18Rik antibody, Actin related protein 2/3 complex inhibitor antibody, Arp2/3 inhibition protein antibody, ARPIN antibody, C15orf38 antibody, Chromosome 15 open reading frame 38 antibody, CO038_HUMAN antibody, Hypothetical protein LOC348110 antibody, MGC61550 antibody, RIKEN cDNA 2610034B18 antibody, UPF0552 protein C15orf38 antibody
Clone:	Polyclonal
Applications:	ELISA WB IHC
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q7Z6K5
Host Species:	Rabbit
Purification:	Antigen Affinity Purified
Immunogen:	Recombinant Human Arpin protein (1-226AA)
Immunogen Species:	Homo sapiens (Human)
Buffer:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Form:	Liquid

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	WB	1:1000-1:5000
	IHC	1:20-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data

Image




Description

Western blot All lanes: ARPIN antibody at 0.5 μ g/ml + HepG2 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 25, 44 kDa Observed band size: 25 kDa



Immunohistochemistry of paraffin-embedded human colon cancer using PACO37454 at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO37454 at dilution of 1:100